#### RESEARCH



# Molecular Features Associated with Resilience to Ocean Acidification in the Northern Quahog, *Mercenaria mercenaria*

 $\label{eq:caroline} Caroline \ Schwaner^1 \cdot \ Sarah \ Farhat^1 \cdot \ Michelle \ Barbosa^1 \cdot \ Isabelle \ Boutet^2 \cdot \ Arnaud \ Tanguy^2 \cdot Emmanuelle \ Pales \ Espinosa^1 \cdot \ Bassem \ Allam^1$ 

Received: 29 September 2022 / Accepted: 8 November 2022 © The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2022

#### Abstract

The increasing concentration of  $CO_2$  in the atmosphere and resulting flux into the oceans will further exacerbate acidification already threatening coastal marine ecosystems. The subsequent alterations in carbonate chemistry can have deleterious impacts on many economically and ecologically important species including the northern quahog (*Mercenaria mercenaria*). The accelerated pace of these changes requires an understanding of how or if species and populations will be able to acclimate or adapt to such swift environmental alterations. Thus far, studies have primarily focused on the physiological effects of ocean acidification (OA) on *M. mercenaria*, including reductions in growth and survival. However, the molecular mechanisms of resilience to OA in this species remains unclear. Clam gametes were fertilized under normal  $pCO_2$  and reared under acidified (pH ~ 7.5,  $pCO_2 \sim 1200$  ppm) or control (pH ~ 7.9,  $pCO_2 \sim 600$  ppm) conditions before sampled at 2 days (larvae), 32 days (postsets), 5 and 10 months (juveniles) and submitted to RNA and DNA sequencing to evaluate alterations in gene expression and genetic variations. Results showed significant shift in gene expression profiles among clams reared in acidified conditions as compared to their respective controls. At 10 months of exposure, significant shifts in allele frequency of single nucleotide polymorphisms (SNPs) were identified. Both approaches highlighted genes coding for proteins related to shell formation, bicarbonate transport, cytoskeleton, immunity/stress, and metabolism, illustrating the role these pathways play in resilience to OA.

Keywords RADSeq · RNASeq · Clam · Ocean acidification · SNPs

| $\bowtie$ | Bassem Allam                |
|-----------|-----------------------------|
|           | bassem.allam@stonybrook.edu |

Caroline Schwaner Caroline.schwaner@stonybrook.edu

Sarah Farhat Sarah.farhat@mnhn.fr

Michelle Barbosa michellebarbosa810@gmail.com

Isabelle Boutet isabelle.boutet@sb-roscoff.fr

Arnaud Tanguy arnaud.tanguy@sb-roscoff.fr

Emmanuelle Pales Espinosa emmanuelle.palesespinosa@stonybrook.edu

<sup>1</sup> School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11790, USA

<sup>2</sup> Station Biologique de Roscoff, CNRS/Sorbonne Université, Place Georges Teissier, 29680 Roscoff, France

# Introduction

The northern quahog (Mercenaria mercenaria) is an ecologically and economically important species within coastal and estuarine ecosystems along the Eastern coast of the United States. As benthic suspension feeders, clams filter algae and suspended particles from the water column which can improve water quality by increasing light penetration for seagrasses (Wall et al. 2008), prevent algal blooms (Cerrato et al. 2004), and contribute to nutrient cycling (Dame 2011). In addition to these beneficial services, M. mercenaria is also one of the most valuable aquaculture species for the east coast and supports both recreational and commercial fisheries. Total clam aquaculture in the US had an estimated value of \$129 million in 2018 (NFMS 2019), and the fishery was valued at \$52.8 million (NMFS 2019). Calcifying marine organisms, such as M. mercenaria, are particularly vulnerable to ocean acidification (OA). Understanding how clams will fare under predicted OA is important as this could have implications for local and national economies and estuarine ecology.

Alterations in carbonate chemistry from elevated  $CO_2$ in seawater will result in a reduction of carbonate ( $CO_3$ ) and calcium carbonate saturation ( $CaCO_3$ ), which will impact marine organisms that biomineralize  $CaCO_3$  to form shells (Feely et al. 2004, 2009; Sabine et al. 2004). Changes to the external seawater chemistry can also affect the acid–base balance of bivalves (Michealidis et al. 2005; Fabry et al. 2008). Evidence suggests that bivalves can respond to OA via adaptive mechanisms including reallocating metabolic resources (Lanning et al. 2010; Thomsen and Melzner 2010; Xu et al. 2016; Gray et al. 2017); however, this can lead to trade-offs between competing physiological processes such as immunity (Burge et al. 2014; Sun et al. 2017; Schwaner et al. 2020).

Some species and populations of marine bivalves are already exposed to coastal waters with low pH and elevated  $pCO_2$  (Wallace et al. 2014). Coastal acidification is a current problem in many estuaries because of excessive nutrient runoff that leads to algal blooms and subsequent microbial decay (Cai et al. 2011; Wallace et al. 2014; Baumann et al. 2015; Laurent et al. 2017). In natural populations of *M. mercenaria*, gonad maturation occurs when water temperatures begin to rise in the late spring and early summer and culminates in spawning when temperatures exceed 20 °C (Rice 1992). Spawning and larval development coincides with coastal waters reaching their extremes of acidification (Wallace et al. 2014; Baumann et al. 2015). Consequently, these animals are already challenged with high CO<sub>2</sub> and low pH during vulnerable stages of development, which will become more extreme with climate change.

Advances in omics have revealed many molecular responses to short term CO<sub>2</sub> stress that might enable organisms to tolerate variable pH (Timmins-Schiffman et al. 2014; Dineshram et al. 2015; Liu et al. 2016; Goncalves et al. 2017; Zhao et al. 2020). Studies of the oysters *Crassostrea gigas* (Timmins-Schiffman et al. 2014), *C. hongkongensis* (Dineshram et al. 2015), *Saccostrea glomerata* (Goncalves et al. 2017), and *Pinctada fucata* (Liu et al. 2016) revealed pathways underlying resilience to OA, including calcium homeostasis, acid base regulation, and metabolism. On the other hand, studies of the transcriptional response of bivalves to OA also reveal depressed pathways (Goncalves et al. 2017; Zhao et al. 2020). For example, the Korean mussel (*Mytilus coruscus*) downregulates gene expression levels of proteins involved in shell formation, resulting in weakened shells (Zhao et al. 2020).

While prior research has primarily focused on acclimation through transcriptomic analyses, studies are beginning to investigate the potential for adaptation to OA by looking at genetic variation and evidence of selection. Differences in the single nucleotide polymorphism (SNP) profiles of the mussel *M. galloprovincialis* between those reared in normal and acidified conditions suggest the potential for rapid adaptation to OA (Bitter et al. 2019). The authors identified hundreds of SNPs responding to each pH treatment and suggested that loci critical for increased fitness under OA are under selection in low pH environments (Bitter et al. 2019). Similarly, studies in the sea urchin *Strongylocentrotus purpuratus* (Pespeni et al. 2013a, b) reported significant allelic changes that suggested the potential for adaptation to OA. In contrast, there was little evidence to support adaptation potential in tropical sea urchin *Echinometra* sp. (Uthicke et al. 2019), highlighting variability between organisms and further demonstrating the need to expand these studies to encompass more species.

Although the physiological impacts of OA on M. mercenaria have been investigated (Talmage and Gobler 2009, 2011; Schwaner et al. 2020), the molecular mechanisms associated with resilience to OA in this species are relatively unknown. The objectives of this study were to determine if OA has a significant impact on the regulation of the transcriptome of M. mercenaria throughout various life history stages and to determine if OA-induced differential mortality allows the identification of SNPs associated with resilience. Here, potential for acclimation and adaptation to OA in M. mercenaria was explored. Impacts of acidification on gene expression and genetic variation throughout development in larvae and juveniles were investigated. Clams were continuously reared in control ( $pCO_2$  of ~ 600 ppm, pH ~ 7.9) or acidified seawater ( $pCO_2$  of ~1200 ppm, pH ~7.5) and samples were collected at 2 days (larvae), 32 days (postsets), 5 months (juveniles), and 10 months (juveniles) postfertilization for analysis of differentially expressed genes (DEGs). Clams that had survived 10 months of chronic OA exposure were deemed as "resilient" to OA, and samples were collected to determine differences in the frequency of SNPs between  $pCO_2$  conditions. Furthermore, the function of differentially expressed genes or genes containing variants was investigated to identify molecular pathways associated with resilience to acidification.

## **Materials and Methods**

#### **Seawater Chemistry and Sample Collection**

In order to increase genetic diversity of experimental animals, wild adult clams from four populations (20 clams from each of the following locations: Wellfleet, Massachusetts (41.9305° N, 70.0310° W); Northport, New York (40.9009° N, 73.3432° W); Hampton Bays, New York (40.8690° N, 72.5176° W); Riverhead, New York (40.9170° N, 72.6620° W) were maintained as broodstock and conditioned for spawning (Helm et al. 2004; Wallace et al. 2008; Schwaner et al. 2020). Females were moved into individual containers for egg collection, and after completion of spawning, eggs and sperm were mixed to ensure genetic heterogeneity. After allowing sufficient time for fertilization (1 h), around 30 million embryos were moved to aquaria with either ambient seawater ( $pCO_2$  of ~ 600 ppm, pH ~ 7.9) or acidified seawater  $(pCO_2 \text{ of} \sim 1200 \text{ ppm}, \text{pH} \sim 7.5)$ . Aquaria were continuously bubbled with 5% CO<sub>2</sub> mixed with air (via multi gas channel proportioners from Cole Parmer (Antylia Scientific; Vernon Hills, IL); Schwaner et al. 2020) or ambient air to maintain  $pCO_2$  conditions. Seawater samples were collected and analyzed (Schwaner et al. 2020) using an EGM-4 Environmental Gas Analyzer ® (PP systems; Amesbury, MA) after acidification and separation of gas phases using a Liqui-cel® Membrane (3 M Membrana; Wuppertal, Germany). pCO<sub>2</sub> total alkalinity, carbonate concentration, and aragonite and calcite saturation were calculated using the R package seacarb (Gattuso et al. 2018), with known first and second dissociation constants of carbonic acid in seawater (Millero 2010). For quality assurance, certified reference materials were analyzed (provided by Andrew Dickson, Scripps Institution of Oceanography) with a 99.99% recovery rate (Tables S1 and S2). Larva rearing followed protocols outlined in previous publications (Helm et al. 2004; Wallace et al. 2008; Schwaner et al. 2020). After 2 days in either control or acidified conditions, about four million larvae from each condition (1 million/replicate) were sampled. Larvae were collected on a 50-micron sieve, rinsed to remove algae and detritus, checked for viability, and the resulting pellet was flash frozen and stored at -80 °C (sampling outlined in Fig. S1). Immediately after settling and metamorphosis (32 days post fertilization, hereby designated "postsets"), 10 juvenile clams (height 0.41 mm  $\pm$  0.1, length 0.44 mm  $\pm$  0.1) per replicate (40 per pCO<sub>2</sub> condition) were flash frozen and preserved at -80 °C. After settling, juvenile clams were transferred from a static system to a flow through system with water from Old Fort Pond in Southampton, NY (40°53'07.2"N 72°26'31.4"W). For the acidified condition, seawater flowed into an acidification chamber where 100% CO<sub>2</sub> was mixed with air via multi gas channel proportions and continuously bubbled to maintain a delta of 0.4 units between the two conditions (carbonate chemistry in Tables S3 and S4). After the seawater reached target values, it flowed into a header tank and the four corresponding replicate tanks using a "downweller" setting. Clams were held on 212 µm nylon mesh sieve and seawater flowed from the top to the bottom compartment of the tank (Schwaner et al. 2020). For the control condition, water flowed into an aerated head tank where it then continuously flowed into each of four replicate tanks. After five months of continuous exposure to either control or acidified conditions in the flow through system, a third sample was collected (10 juveniles/ replicate,  $40/pCO_2$  condition; length 3.65 mm  $\pm$  1.21; height  $3.34 \pm 1.07$ ). Then after 10 months, samples for both RNA sequencing (10 juveniles/replicate, 40/pCO<sub>2</sub> condition) and DNA sequencing (12 juveniles/replicate, 48/pCO<sub>2</sub> condition, 96 total) were collected (length 5.50 mm  $\pm$  1.39; height

4.86 mm  $\pm$  1.23). Before DNA extraction, juvenile clam tissues were removed from the shells and added to 100% molecular grade ethanol.

#### **RNA and DNA Extraction**

RNA was extracted from samples collected at 2 and 32 days, 5 and 10 months using NucleoSpin® RNA Plus Kit (#740,984.50; Macherey-Nagel; Düren, Germany) where DNA was removed with DNA-free<sup>TM</sup> Kit (#AM1906; Ambion, Inc.; Austin, TX), following manufacturer's protocols. For larval samples, RNA was extracted from a pellet of ~ 100,000 larvae, and for juveniles RNA was extracted from individual whole bodies. RNA quality and quantity were checked using a NanoDrop® ND-1000 Spectrophotometer (Thermo Fisher Scientific; Waltham, MA). At 10 months, samples were also collected for DNA Sequencing. DNA was isolated from samples using standard phenol:chloroform:isoamyl extraction (Farhat et al. 2020; Boutet et al. 2022). DNA quality and quantity were checked using the Qubit dsDNA HS Assay Kit with assays read on a Qubit 2.0 fluorometer (Life Technologies/Agilent 2100; Waldbronn, Germany) and by agarose gel electrophoresis.

# RNAseq Library Preparation, Sequencing, and Analysis

Extracted RNA was sent to Novogene Corporation (Sacramento, CA, USA) for sequencing. Sequencing libraries were generated using NEBNext® Ultra<sup>™</sup> RNA Library Prep Kit for Illumina® (# E7760L; New England Biolabs; Ipswich, MA), following manufacturer's instructions. Libraries were sequenced on Illumina platform (Novaseq 2X150) and 150 paired-end length reads were then demultiplexed. The final reads received were inputted into the CLC workbench (version 11.0.1 (https://digitalinsights.qiagen.com), and trimmed based on quality scores (limit 0.05), ambiguous nucleotides (max 2 nucleotides per sequence), and adapters. Trimmed reads were mapped onto the M. mercenaria gene prediction (Farhat et al. 2022) using the "map reads to reference" function with default parameters. The next in-silico analyses were performed on the computing cluster SeaWulf at Stony Brook University. Reads were sorted, indexed, and the number of reads mapping to transcripts was quantified using Samtools version 1.9 (Li 2011). Counts were normalized and compared between conditions at various times of exposure (2 and 32 days, 5 and 10 months) using DESeq2 (1.36.0) from Bioconductor (Love et al. 2014). The p-values were corrected for multiple testing using the Benjamini and Hochberg method. DEGs with a log fold change >|2| and adjusted p-value < 0.05 were deemed significant. Principal component analysis (PCA) was performed using plot-PCA function in DE Seq2 of the vsd-transformed (variance stabilizing transformation) counts, volcano maps were created to visualize fold change (Fig. S2), and cluster heat maps were generated for the top DEGs (61-100 depending on number of DEGs) (Fig. S3). We used the functional annotation of *M. mercenaria* proteins generated from the gene annotation (Farthat et al. 2022). A hypergeometric test was performed to test for statistical enrichment (FDR adjusted p value < 0.05) of GO terms using the *TopGO* package in R.

# ddRAD Library Preparation, Sequencing, and Analysis

ddRAD libraries were prepared using a protocol with steps modified from Peterson et al. (2012). Briefly, 200 ng of each DNA sample was double digested with PstI high fidelity restriction endonuclease and MseI (New England Biolabs). After 10 h at 37 °C, the enzymes were inactivated by heating at 65 °C for 20 min. DNA samples were individually identified through the ligation of specific P1-n barcoded adapters at 16 °C for 6 h in a 30 µl reaction containing P1 adapter, T4 DNA ligase, 1 mM ATP, P2 adapter, and 1X CutSmart reaction buffer (New England Biolabs). Samples were then purified and size selected using Nucleomag NGS clean-up and size select kit (#744,970.50; Machery-Nagel) at 1:1 ratio of beads to sample and then resuspended in Tris buffer. Purified adapter-ligated fragments were then PCR-amplified with 12 uniquely indexed PCR-P2 primer sequences for multiplexing, allowing for post-sequencing demultiplexing. Gel electrophoresis was performed on PCR products and visualized using GelRed Stain and additional PCR was run if necessary. Each PCR product was then quantified using Qubit dsDNA HS assay kit, and then size selected with Pippin-Prep (1.5% agarose dye free gel cassette, Sage Science; Beverly, MA) under a setting with a range of 350-800 bp. Each fraction (350-800 bp, and 380-800 bp) was quantified with Qubit and quality was checked on a Bioanalyzer with high sensitivity DNA chip (Agilent technologies). To have better distribution, a larger range of sizes, and better representativity of each size fraction we added 50% of the 350-800 bp, and 50% of the 380–800 bp cut to the library for sequencing. 5% of PhiX control libraries were used to create a more diverse set of clusters as is recommended by Illumina for our sample type. Samples were sequenced using Illumina NovaSeq 6000 System with an S4 flow cell (Dr. Graham Wiley, Oklahoma Medical Research Foundation) on one lane.

The resulting paired end reads were demultiplexed by index by Novogene and then demultiplexed by barcode into individual samples using CLC Workbench. Reads were trimmed as described above and then mapped onto the *M. mercenaria* reference genome (Farhat et al. 2022) using CLC "map reads to reference" function with default parameters. This was followed by running the "Local Realignment" tool (selecting to realign unaligned ends with 3 multi passes using guiding variants). Bam files were exported from CLC to Stony Brook's high performance computing cluster, SeaWulf (Stony Brook University 2016) and variants were identified using Stacks v2 with the ref\_map pipeline (Rochette et al. 2019). Data filtering for detection of high-quality SNPs included: minimum allele frequency 0.05 (-min-maf), a minimum percentage of samples per population (75%), a HWE p-value of < 0.05 (-hwe) (the probability that this variant deviates from HWE). After filtering, only SNPs that had significantly higher F<sub>ST</sub> between the compared populations values were selected (corrected AMOVA  $F_{ST} > 0.05$  and p < 0.05). The position of each variant in the genome (intergenic, untranslated region, coding sequence, or intron) and the type of consequence the variant has on the resulting protein (missense, synonymous, stop gained) were inferred as described by Farhat et al. (2020). To visualize population structure, we performed a principal components analysis to see variation between samples, using the *glPCA* function in R.

# Results

We firstly tested the hypothesis that OA induces a significant regulation of the M. mercenaria transcriptome and identified genes that are differentially regulated in response to OA. At all life history stages investigated 2 day (larvae), 32 days (postsets), 5 months (juveniles), 10 months (juveniles) there were DEGs between clams in acidified (pH 7.5) and control conditions (pH 7.9). In addition, we explored the possibility that OA leads to selection that can be detected in the SNP profile of survivors and that certain genotypes are resilient to OA. Genetic difference between acidified and control clams at 10 months was overall low; however, there was significant enrichment of SNPs in clams surviving acidified conditions. The significant SNPs and DEGs were found in genes coding for proteins involved in biological processes known to be impacted by OA. These included genes related to shell formation, bicarbonate transport, cytoskeleton, immunity/ stress, and metabolism, highlighting the role these pathways play in resilience to OA. The following sections provide thorough results from gene expression and SNP identification for clams under acidification stress.

#### **Differentially Expressed Genes**

Gene expression analysis revealed variations in the transcriptome of clams reared in acidified and control conditions in every life stage investigated. While there was evident variation from  $pCO_2$ , there were also differences between the developmental stages. The fewest DEGs were found in larvae that were exposed to acidified conditions for only 2 days and juveniles in acidified conditions for 10 months. T he **Fig. 1** Differentially expressed gene counts of *M. mercenaria* at 2, 32 days, 5 and 10 months of OA exposure. Number of genes upregulated under acidification stress (shades of red) and downregulated (shades of blue) at each sampling time point



greatest number of DEGs were detected in the 5 month old juvenile clams (Fig. 1). In the postset juveniles (32 days), 92% of their DEGs were downregulated in the acidified condition. Similarly, for 10 month old juveniles 80% of the DEGs were downregulated in the acidified condition.

PCA showed clear separation between  $pCO_2$  treatments, with the 32 day old post set juveniles having the greatest separation between control and acidified clams (Fig. 2). PCA resulted in clustering of samples by age, showing distinct segregation between larval and juvenile samples and the most overlap between 5 and 10 months (Fig. 2).

Several genes upregulated under acidified conditions were similarly expressed between the different life stages (Fig. 3). In addition, many genes upregulated under acidified conditions coding for genes with the same predicted protein, were similarly expressed between life stages (Fig. 4). These genes and proteins were primarily involved with biomineralization and acid-base regulation.



**Fig. 2** Principal component analysis (PCA) of normalized gene expression data at 2 days, 32 days, 5 months, and 10 months between clams grown under different  $pCO_2$  conditions (red acidified; blue control)

**Fig. 3 A** The Venn diagram presents the number of genes uniquely expressed within each group with the overlapping regions showing the number of genes that are expressed in 2 or more groups **B** DEGs in common/shared between comparisons, \* is next to group with higher expression for the gene in acidified conditions. Life stages are 2d (2 days), 32d (32 days) 5 m (5 months) and 10 m (10 months)



# В

| Gene ID                   | Protein  | Life stage |
|---------------------------|--|------------|
| mRNA.chromosome_12.2645.1 | Uncharacterized  | 2*, 5*     |
| mRNA.chromosome_6.995.1   | VWFC domain  | 2*, 5*     |
| mRNA.chromosome_5.710.3   | sterol regulatory element-binding transcription factor 1 | 2*,5*      |
| mRNA.chromosome_12.282.1  | Chitin-binding type-2 domain profile                     | 2*, 5*     |
| mRNA.contig_3016.1.1      | Uncharacterized  | 2, 32      |
| mRNA.chromosome_12.297.1  | Collagen, type XII, alpha                                | 32*, 10    |
| mRNA.chromosome_1.1199.2  | MFS transporter, OCT family, solute carrier family 22    | 32*, 10    |
| mRNA.chromosome_12.2684.1 | Histone H4   | 32, 5      |
| mRNA.chromosome_12.2682.1 | Histone H4   | 32, 5*     |

These included genes with Von Willebrand Factor (VWF) domains, chitin-binding domains, epidermal growth factor (EGF) domains, and solute carriers (Figs. 3 and 4).

## Two Day Old Larvae

Genes coding for proteins related to the cytoskeleton, shell formation, protein binding, and bicarbonate transport were upregulated in two day old larvae in the acidified condition (Tables 1, 2 and S5). Moreover, chitin metabolic process was a significantly enriched GO term within the upregulated genes (Table S6). Transcription related genes (such as gene with Zinc finger CCHC-type profile and aryl hydrocarbon receptor nuclear translocator) were downregulated and transcription factor complex was the only significantly enriched GO term in downregulated genes.

#### 32 Day Old Juveniles

Postset juveniles had almost twice as many DEGs as larvae (Table 1), with the majority downregulated. Upregulated genes included those related to shell formation, bicarbonate transport, and immune/stress processes (Tables 3 and S5). Similar to 2 day old larvae, chitin metabolic process was also a significantly enriched GO term within upregulated genes (Table S7). Genes related to cytoskeleton and immune/stress processes were downregulated (Table 3).

## **Five Month Old Juveniles**

After five months of exposure to OA, the juveniles had the greatest number of DEGs and the greatest percentage upregulated (Table 1). Upregulated genes included those related to cytoskeleton, shell formation, bicarbonate

Fig. 4 A The Venn diagram presents the number of genes coding for unique proteins within each group with the overlapping regions showing the number of genes coding for similar proteins that are expressed in 2 or more groups **B** DEGs coding for proteins in common/shared between comparisons, \* is next to group with higher expression for the gene in acidified conditions, the number of genes having the subject domain, including those upregulated under acidified conditions, is given between parentheses when more than one is found in that life stage



# В

| Protein/Domain                    | Life stage             |
|-----------------------------------|------------------------|
| Solute carrier                    | 2*, 32*, 5 (4-2up), 10 |
| Chitin binding; Chitin synthase   | 2*, 32*, 5(3)*, 10     |
| EGF                               | 2*, 32*, 5(2)*         |
| Zinc Finger CCHC                  | 2 (4-1up), 5(2-1up),   |
| VWF                               | 2(2)*,32(2)*,5(5-3up)  |
| Sterol regulatory binding element | 2*,5*                  |
| GTP binding protein               | 2*,10*                 |
| Vitelline                         | 32*, 5*,10             |
| Collagen                          | 32*,10                 |
| Cathepsin                         | 32*,5(3)*              |
| Histone h4                        | 32(2),5(2-1up)         |
| G protein                         | 32,5*                  |
| Serine/threonine-protein          | 5*, 10*                |
| TNF                               | 5*, 10                 |
| Cytochrome p450                   | 5*, 10                 |

| Table 1  | Number of total DEGs |
|----------|----------------------|
| between  | comparisons and      |
| percent  | of DEGs up or down   |
| regulate | d                    |

|                        | DEGs | Upregulated | Percent<br>upregulated | Downregulated | Percent<br>downregulated |
|------------------------|------|-------------|------------------------|---------------|--------------------------|
| 2 day old larvae       | 61   | 26          | 42.6%                  | 35            | 57.4%                    |
| 32 day old post sets   | 108  | 9           | 83.3%                  | 99            | 91.7%                    |
| 5 month old juveniles  | 170  | 112         | 65.9%                  | 58            | 34.1%                    |
| 10 month old juveniles | 20   | 4           | 20%                    | 16            | 80%                      |

| ID                        | Predicted Protein                            | Log2FC | Padj | Functional Group      |
|---------------------------|--|--------|------|-----------------------|
| mRNA.chromosome_19.1142.1 | Coadhesin                                    | 2.7    | 0.05 | Cytoskeleton          |
| mRNA.chromosome_17.1806.1 | Putative GTP-binding protein6                | 2.2    | 0.02 | Cytoskeleton          |
| mRNA.chromosome_12.824.1  | Fibronectin type-III domain profile          | 2.4    | 0.04 | Shell formation       |
| mRNA.chromosome_6.995.1   | VWFC domain profile                          | 2.4    | 0.00 | Shell formation       |
| mRNA.chromosome_12.1499.3 | Aggrecan 1 protein                           | 2.0    | 0.03 | Shell formation       |
| mRNA.chromosome_12.282.1  | Chitin-binding type-2 domain profile         | 2.0    | 0.02 | Shell formation       |
| mRNA.chromosome_14.664.1  | Calcium-binding EGF-like domain              | 2.0    | 0.03 | Shell formation       |
| mRNA.chromosome_3.211.2   | VWFD domain profile                          | 2.0    | 0.01 | Shell formation       |
| mRNA.chromosome_15.2198.2 | F-box only protein 38 isoform X2             | 2.3    | 0.03 | Protein binding       |
| mRNA.chromosome_5.62.1    | Translation initiation factor 5B             | 2.3    | 0.02 | Protein binding       |
| mRNA.chromosome_1.3867.1  | RNA-binding protein 25                       | 2.7    | 0.03 | RNA binding           |
| mRNA.chromosome_8.323.1   | Zinc finger CCHC domain-containing protein 9 | 2.3    | 0.03 | RNA binding           |
| mRNA.chromosome_10.427.1  | Solute carrier family 34                     | 3.6    | 0.00 | Bicarbonate transport |

**Table 2** Representative significantly upregulated genes in acidified larvae. Genes presented here were selected based on log2foldchange>2,adjusted p-value < 0.05, and relevance of function

transport, and immunity (Tables 4 and S5). Transmembrane transporter activity and transporter activity were significantly enriched GO terms among upregulated genes (Table S8). Downregulated genes in juvenile clams included those related to cytoskeleton, ion transport, metabolism, and immune/stress processes (Table 4). GO terms of downregulated genes were enriched for metabolic processes (Table S8).

#### **10 Month Old Juveniles**

The number of DEGs of 10 month old juveniles was the fewest with the majority downregulated. Upregulated genes in 10 month old clams exposed to OA included those related to the cytoskeleton, DNA and nucleotide repair, and RNA binding. (Tables 5 and S5). The downregulated genes were related to the shell formation, immune/stress processes,

**Table 3** Representative significantly upregulated genes in acidified postset juveniles. Genes presented here were selected based on log2fold-<br/>change > 2, adjusted p-value < 0.05, and relevance of function

| ID                        | Predicted Protein  | Log2FC | Padj    | Functional Group      |
|---------------------------|--|--------|---------|-----------------------|
| mRNA.chromosome_12.297.1  | Collagen, type XII, alpha  | 6.6    | 0.01    | Shell formation       |
| mRNA.chromosome_14.667.1  | EGF-like domain profile  | 6.0    | 0.02    | Shell formation       |
| mRNA.chromosome_15.1602.1 | WAP-type 'four-disulfide core' domain profile                        | 5.6    | < 0.001 | Shell formation       |
| mRNA.chromosome_15.1544.1 | Von Willebrand factor type A domain                                  | 5.7    | 0.01    | Shell formation       |
| mRNA.chromosome_7.1094.1  | Chitin-binding protein   | 4.6    | 0.02    | Shell formation       |
| mRNA.chromosome_13.1301.1 | Von Willebrand factor type C domain                                  | 2.0    | < 0.001 | Shell formation       |
| mRNA.chromosome_1.1199.2  | MFS transporter, OCT family, solute carrier family 22                | 3.0    | 0.02    | Bicarbonate transport |
| mRNA.chromosome_15.882.1  | Vitelline membrane outer layer protein 1                             | 6.7    | < 0.001 | Immune                |
| mRNA.chromosome_10.766.1  | Heat shock protein 70 kDa protein 12A like                           | -4.8   | < 0.001 | Stress/immunity       |
| mRNA.chromosome_9.410.1   | Acidic leucine-rich nuclear phosphoprotein 32 family<br>member A/C/D | -4.2   | < 0.001 | Stress/immunity       |
| mRNA.chromosome_13.2893.1 | Heat shock protein 70  | -3.5   | 0.002   | Stress/immunity       |
| mRNA.chromosome_12.2682.1 | Histone H4   | -3.2   | 0.001   | Stress/immunity       |
| mRNA.chromosome_12.2684.1 | Histone H4   | -3.0   | < 0.001 | Stress/immunity       |
| mRNA.chromosome_2883.2.1  | Mucin 5B   | -4.6   | < 0.001 | Cytoskeleton          |
| mRNA.chromosome_5.2802.1  | tRNA modification GTPase   | -4.0   | < 0.001 | Cytoskeleton          |
| mRNA.chromosome_19.81.1   | Rem2- and rab-like small GTPase 1                                    | -3.7   | < 0.001 | Cytoskeleton          |
| mRNA.chromosome_3797.3.1  | Rem2- and rab-like small GTPase 1                                    | -3.1   | < 0.001 | Cytoskeleton          |
| mRNA.chromosome_6.871.2   | Tubulin polyglutamylase TTLL2  | -2.7   | < 0.001 | Cytoskeleton          |

| ID                        | Predicted Protein                             | Log2FC | Padj    | Functional Group      |
|---------------------------|---|--------|---------|-----------------------|
| mRNA.chromosome_12.726.1  | Actin beta/gamma 1                            | 6.7    | < 0.001 | Cytoskeleton          |
| mRNA.chromosome_1.2647.1  | omosome_1.2647.1 Dynein light chain           |        | 0.00    | Cytoskeleton          |
| mRNA.chromosome_5.1141.1  | Myosin heavy chain                            | 5.3    | 0.00    | Cytoskeleton          |
| mRNA.chromosome_3.1998.1  | Actin beta/gamma 1                            | 5.9    | 0.00    | Cytoskeleton          |
| mRNA.chromosome_2.2738.1  | Actin beta/gamma 1                            | 4.4    | 0.01    | Cytoskeleton          |
| mRNA.chromosome_13.1054.1 | Tubulin monoglycylase TTLL3/8                 | 3.1    | 0.01    | Cytoskeleton          |
| mRNA.chromosome_11.1042.1 | Actin beta/gamma 1                            | 2.5    | 0.01    | Cytoskeleton          |
| mRNA.chromosome_13.2773.1 | Chitin-binding type-2 domain profile          | 6.9    | < 0.001 | Shell formation       |
| mRNA.chromosome_4.1315.1  | Chitin-binding type-2 domain profile          | 6.7    | < 0.001 | Shell formation       |
| mRNA.chromosome_1.1197.1  | EGF-like signature                            | 6.4    | 0.00    | Shell formation       |
| mRNA.chromosome_6.995.1   | VWFC domain profile                           | 6.4    | 0.00    | Shell formation       |
| mRNA.chromosome_7.1323.1  | EGF domain                                    | 6.1    | 0.00    | Shell formation       |
| mRNA.chromosome_12.282.1  | Chitin-binding type-2 domain profile          | 6.0    | 0.00    | Shell formation       |
| mRNA.chromosome_3.2434.1  | Von Willebrand factor type A domains          | 5.8    | 0.00    | Shell formation       |
| mRNA.chromosome_9.961.1   | EF-hand calcium-binding domain profile        | 5.7    | 0.00    | Shell formation       |
| mRNA.chromosome_6.581.1   | EF-hand calcium-binding domain                | 5.2    | 0.05    | Shell formation       |
| mRNA.chromosome_5.1518.1  | Phosphorylase kinase alpha/beta subunit       | 3.6    | 0.01    | Shell formation       |
| mRNA.chromosome_6.997.1   | Von Willebrand factor type A domains          | 3.3    | 0.01    | Shell formation       |
| mRNA.chromosome_13.1112.1 | C-type lectin -type domain                    | 3.0    | 0.01    | Shell formation       |
| mRNA.chromosome_1.284.1   | Calmodulin                                    | 2.7    | 00.01   | Shell formation       |
| mRNA.chromosome_15.1517.1 | Solute carrier family 16                      | 6.1    | 0.00    | Bicarbonate transport |
| mRNA.chromosome_7.136.1   | Solute carrier family 16                      | 6.3    | < 0.001 | Bicarbonate transport |
| mRNA.chromosome_9.1391.1  | Vitelline membrane outer layer protein 1      | 6.7    | < 0.001 | Immune                |
| mRNA.chromosome_13.1968.1 | Galectin                                      | 6.3    | 0.00    | Immune                |
| mRNA.chromosome_4.788.1   | Baculoviral IAP repeat-containing protein 2/3 | 6.3    | 0.00    | Immune                |
| mRNA.chromosome_11.833.1  | Phage lysozyme 2                              | 6.2    | 0.00    | Immune                |
| mRNA.chromosome_12.2682.1 | Histone                                       | 5.7    | 0.00    | Immune                |

**Table 4** Representative significantly upregulated genes in 5 month old juveniles. Genes presented here were selected based on  $\log 2$  fold-<br/>change > 2, adjusted p-value < 0.05, and relevance of function</th>

metabolism, DNA binding, and bicarbonate transport (Table 5). GTP binding was an enriched GO term among upregulated genes (Table S9).

# **Genetic Variation**

Mapped reads for each clam averaged 4,125,916  $\pm$  973,563 with 91% mapping to reference genome for controls and 4,714,029  $\pm$  1,026,685 with 95% mapping to the reference genome for acidified clams. Survivors of 10 months of exposure to acidified conditions were contrasted with clams maintained in control conditions. SNPs were identified in each group and filtered for significant variants (Table S10). The F<sub>ST</sub> value between the control and acidified groups was 0.008 (Table S11) and Phi<sub>ST</sub> was 0.003 (Table S12) and PCA showed little difference between pH treatments except in the lower two quartiles.. Summary genetic statistics of each group (acidified or control) based on the entire set of SNPs are summarized in Table S12.

The position of each SNP in the different regions of the genome (coding sequences, untranslated region, intron, intergenic region) was identified (Table S13). Variants in CDS regions represented 4.3% of the total variants, representing about twice as much as CDS coverage in the genome (Farhat et al. 2022). Predicted proteins of significant variants are listed in Table S10 and the consequences of the variants (missense, stop-gained, synonymous) detected in CDS are given in Table S14. We identified 1,668 variants in the CDS regions, including 398 missense, 718 stop-gained, and 552 synonymous variants (Table S14). There were three unique SNPs (2 missense and 1 synonymous) that were only detected in the acidified population and none were unique to control oysters.

SNPs in 233 genes were found with a higher frequency in clams surviving the acidified condition (Table S10). These genes were related to cytoskeleton, calcium binding, shell formation, growth/development, immune/stress, metabolism, signaling, protein synthesis, and bicarbonate transport (Tables 6, S10). Only three of these SNPs were

| ID                        | Predicted Protein  | Log2FC | padj | Functional Group             |
|---------------------------|--|--------|------|------------------------------|
| mRNA.chromosome_4.841.1   | Serine/threonine-protein phosphatase 6 regulatory ankyrin repeat subunit A |        | 0.05 | DNA repair and recombination |
| mRNA.chromosome_7.1744.2  | Cullin 4   | 8.02   | 0.01 | Nucleotide excision repair   |
| mRNA.contig_3511.13.1     | Cleavage stimulation factor subunit 1-like                                 | 7.87   | 0.03 | RNA binding                  |
| mRNA.chromosome_17.349.1  | ADP-ribosylation factor 6-like   | 7.10   | 0.05 | Cytoskeleton                 |
| mRNA.chromosome_11.487.1  | Cytochrome P450, family 4, subfamily B, polypeptide 1                      | -7.24  | 0.02 | Immune/stress                |
| mRNA.chromosome_16.585.1  | Vitelline membrane outer layer protein 1 homolog                           | -7.47  | 0.01 | Immune/stress                |
| mRNA.contig_3987.2.1      | Plasminogen  | -6.91  | 0.05 | Immune/stress                |
| mRNA.contig_3781.1.1      | TNF(Tumour Necrosis Factor) family   | -6.82  | 0.05 | Immune/stress                |
| mRNA.chromosome_15.1640.2 | Insoluble matrix shell protein 5-like                                      | -8.20  | 0.05 | Shell formation              |
| mRNA.chromosome_12.297.1  | Collagen, type XII, alpha  | -7.06  | 0.03 | Shell formation              |
| mRNA.chromosome_1.1199.2  | Solute carrier family 22   | -7.58  | 0.01 | Bicarbonate transport        |
| mRNA.chromosome_5.3241.1  | Dopamine beta-monooxygenase; temptin-like isoform X1                       | -3.78  | 0.05 | Metabolism                   |
| mRNA.chromosome_9.1656.1  | Fibroblast growth factor receptor 1  | -6.95  | 0.05 | Growth factor                |
| mRNA.chromosome_11.2178.1 | Serrate RNA effector molecule homolog isoform X2                           | -7.74  | 0.01 | DNA binding                  |
| mRNA.chromosome_1.412.1   | Serrate RNA effector molecule homolog isoform X2                           | -7.64  | 0.01 | DNA binding                  |
| mRNA.chromosome_2.93.1    | Serine/arginine repetitive matrix protein 5-like isoform X2                | -6.88  | 0.05 | DNA binding                  |

**Table 5** Representative significantly downregulated genes in 10 month old juveniles. Genes presented here were selected based on log2fold-change > 2, adjusted p-value < 0.05, and relevance of function.</th>

unique (absent in control clams). These included a missense SNP in triple functional domain protein, a missense SNP in relaxin family peptide receptor 1, and a synonymous SNP in an uncharacterized gene. Genes containing SNPs had predicted proteins that were the same as DEGs upregulated under acidification (Table 6). These included a SNP in the same exact gene "mRNA.chromosome\_17.1526.1" which codes for adapter protein CIKS-like. Overrepresented GO terms of SNPs significantly enriched in acidified clams are listed in Table S15.

SNPs in 222 genes were found to be depleted in clams reared in acidified conditions (Table S10). None of the SNPs depleted in acidified clams were unique to the control population. In general, the SNPs were found in genes with the same broad functional groups as those enriched in acidified conditions (Table S10). GO terms enriched in depleted genes are listed in Table S15.

# Discussion

#### Life History Stage and Length of pCO<sub>2</sub> Exposure

Larvae collected after 2 days of OA exposure had a surprisingly low number of DEGs when compared to the postset and juvenile clam samples. This was unexpected because the first 48 h of larval development is associated with increased vulnerability to OA (Waldbusser et al. 2013). A potential reason for the low number of DEGs might be because early larval development includes many short-lived ontogenetic stages, characterized by unique gene expression. For example, Lopez-Landavery et al. (2021) found differences in gene expression between control geoduck (Panopea globosa) larvae and acidified larva, but with differences between days. At two days, M. mercenaria should all be at the fully shelled D-stage; however, many studies have demonstrated delays in development under OA conditions (Waldbusser et al. 2015; Gray et al. 2017; De Wit et al. 2018). Because larval samples were pooled, there might have been intrinsic variations related to staggered developmental stages within each  $pCO_2$ treatment that lowered the signal from pH treatments themselves. In fact, the surviving acidified larvae were actually larger than the controls five days post fertilization (Schwaner et al. 2020), suggesting that the smaller size cohort may not have survived. This cohort was sampled in the two day time point and could have contributed to variability within the  $pCO_2$  treatment. Despite the lower-than-expected number of DEGs, this information is still useful in understanding the molecular mechanisms of resilience to OA in larval clams, particularly since molecular responses to OA can be species-specific. Upregulated DEGs included genes coding for proteins related to the cytoskeleton, shell formation, and bicarbonate, suggesting an upregulation of cellular growth and calcification to form the fully shelled D-stage larvae despite the alterations in carbonate chemistry of the external seawater.

The second sampling point immediately followed metamorphosis, one of the most dramatic physiological processes during bivalve development (Garcia-Esquivel et al. 2001; Miller et al. 2009). Metamorphosis is energetically

| Table 6 | SNPs in representative | genes with h | nigher freque | encies in acidif | ied clams. | * indicates si | imilar predi | cted protein | as a previously | identified |
|---------|------------------------|--------------|---------------|------------------|------------|----------------|--------------|--------------|-----------------|------------|
| DEG.    |                        |              |               |                  |            |                |              |              |                 |            |

| ID   | Predicted Protein  | SNP   |
|--|--|---|
| mRNA.chromosome_14.727.1   | VWFA domain profile*   | Missense  |
| mRNA.chromosome_1.3056.1   | VWF type C domain*   | Stop gained                                       |
| mRNA.chromosome_19.1514.1; mRNA.chromo-<br>some_2.361.1  | Calcium-binding EGF-like domain signature*   | Missense; stop gained                             |
| mRNA.chromosome_10.1978.1; mRNA.chromo-<br>some_18.1515.1  | Multiple EGF-like domain protein 11;6  | Stop gained; Synonymous                           |
| mRNA.chromosome_1.1227.1   | EGF-like domain profile*   | Synonymous  |
| mRNA.chromosome_13.2030.2  | Cadherin EGF LAG seven-pass G-type receptor 1  | Stop gained                                       |
| mRNA.chromosome_15.2234.2  | Laminin EGF domain   | Missense  |
| mRNA.chromosome_8.1485.1   | Chitin synthase  | Missense  |
| mRNA.chromosome_18.845.1   | Chitin-binding type-2 domain profile*  | Stop gained                                       |
| mRNA.chromosome_14.668.1   | WAP-type (Whey Acidic Protein) 'four-disulfide core'*  | Missense  |
| mRNA.chromosome_2.2663.1; mRNA.chromo-<br>some_5.1350.1  | Protocadherin Fat 1/2/3; protocadherin delta 1   | Stop gained; synonymous                           |
| mRNA.chromosome_5.349.1  | Receptor-type tyrosine-protein phosphatase beta  | Missense  |
| mRNA.chromosome_15.2303.1  | Collagen alpha-6(VI) chain *   | Synonymous  |
| mRNA.chromosome_18.87.1  | IQ calmodulin-binding motif  | Stop gained                                       |
| mRNA.chromosome_16.372.1   | Vitamin D3 receptor-like isoform   | Synonymous  |
| mRNA.chromosome_2.611.1; mRNA.chromo-<br>some_7.210.1; mRNA.contig_4409.2.1                                      | MFS transporter, LAT3 family, solute carrier family 43,<br>member; solute carrier family 16 member 9; solute car-<br>rier family 12 member 8-like* | Stop gained                                       |
| mRNA.chromosome_1.3787.1   | GTP-binding protein 2-like*  | Synonymous  |
| mRNA.chromosome_4.1477.1   | Baculoviral IAP repeat-containing protein 2/3*   | Synonymous  |
| mRNA.chromosome_10.55.1  | Ankyrin  | Synonymous  |
| mRNA.chromosome_12.2299.1; mRNA.chromo-<br>some_12.2378.1; mRNA.chromosome_1.3528.1;<br>mRNA.chromosome_5.2557.1 | Tripartite motif-containing protein 71; 2/3;24; 2/3  | Stop gained; stop gained;<br>synonymous; missense |
| mRNA.chromosome_3.830.1; mRNA.chromo-<br>some_3.1067.1   | Apolipoprotein B   | Stop gained                                       |
| mRNA.chromosome_1.1767.1   | NF-kappa-B inhibitor alpha   | Missense  |
| mRNA.chromosome_18.733.1   | Nuclear factor interleukin-3-regulated protein-like  | Missense  |
| mRNA.chromosome_7.466.1  | Interleukin-1 receptor-associated kinase 4   | Missense  |

demanding, resulting in clams with a reduced energy budget (Videla et al. 1998; Garcia-Esquivel et al. 2001). There was a general downregulation of gene expression in postsets from acidified conditions with over 90% of DEGs with significantly lower expression than controls. This was the greatest percentage of downregulated genes, and PCA revealed that the postsets had the greatest spatial separation from all other time points and between  $pCO_2$  treatments. A general trend of downregulation of gene expression under OA conditions is not surprising. Sea urchin larvae (Lytechinus picitus) exposed to high  $pCO_2$  had significantly different gene expression than sea urchins maintained in control conditions, with 80% of the genes with the largest changes in gene expression being downregulated. These genes were involved in energy supply and demand, which suggests metabolic depression (O'Donnell et al. 2010). Many studies have shown that it is energetically costly to develop under OA (Timmins-Schiffman et al. 2012; White et al. 2013), and differing allocation of energy reserves can result in delayed development and metamorphosis (Talmage and Gobler 2009; Timmins-Schiffman et al. 2012; Wessel et al. 2018). Impacts from OA incurred during the larval stage, such as depletion of energy reserves, can have carry over effects in later juvenile stages, possibly as a result of epigenetic modifications. In the Olympia oyster (Ostrea lurida) impacts from OA during the larval stage lasted 1.5 months after juveniles were transferred to a common environment (Hettinger et al. 2012). The larval clams from acidified conditions overexpressed many genes associated with biomineralization resulting in significantly larger larvae. However, this trend reversed at the juvenile stage, with significantly smaller postset clams from the acidified conditions (Schwaner et al. 2020). This might suggest depleted energy reserves at the larval stage which carried over into the juvenile stage and manifested in reduced growth.

Contrary to the general downregulation of DEGs at 32 days, the five month old juvenile clams had the greatest number and percentage of upregulated DEGs. This sampling point was right before winter, when food would be abundant and temperatures warm enough to increase metabolic activity. During winter, clams are less active, pump less, and deplete energy stores for metabolism. When water temperatures increased in the spring, food levels would still be low (there was no supplemental feeding on the flow through) and metabolic demand would be high, resulting in clams with reduced physiological condition (Zarnoch and Schreibam 2008). The 10 month old clams had 80% of their DEGs downregulated and had less clustering by pCO<sub>2</sub> treatment from PCA than samples from the other time points. The added stress of overwintering might be the cause for this decrease in DEGs and general trend of downregulation as they need to conserve more energy. In addition, studies have demonstrated a threshold for upregulating genes under stressful environmental conditions. For example, heat shock protein 70 was upregulated under OA in the starfish Asterias rubens; however, it was reduced after long term exposure (Henroth et al. 2011). This might suggest it is too costly to overexpress genes for extended periods and that the clams may have reached their threshold sometime between fiveand 10 months. The 10 month old clams also downregulated genes related to shell formation, immunity, and bicarbonate transport, potentially indicating costs of chronic exposure to OA. Alternatively, by this point there might have been a selection of resilient genotypes rather than differential gene expression.

#### **Genetic Differentiation**

Previous research has demonstrated that OA significantly increases mortality of M. mercenaria larvae (Talmage and Gobler 2010, 2011) suggesting that there could be a selection of "OA-resilient" clams. Selection of OA-resilient bivalves has been found in other species (Thomsen et al. 2017; Bitter et al. 2019). Here, there were differences in allele frequencies between clams from control and acidified conditions; however, genetic differences between controls and survivors of OA were relatively low. There were also no unique SNPs in the control treatments and only three (2 missense and 1 synonymous) in the acidified condition. Visualization of SNP data showed clustering, but not by pCO<sub>2</sub>. Four parent populations were spawned to produce the pooled larvae used here. A study of clam population structure along the east coast of North America showed genetically distinct populations between Massachusetts and Mid-Atlantic (New York) populations (Ropp 2020). Within each of the four groups, there appeared to be separation by  $pCO_2$  treatment; however, the genetic structure from source population may have overshadowed the effect of  $pCO_2$  treatment.

Two months after the last sampling point (1 year post fertilization), a reciprocal transplant was performed to test recovery potential and acclimation to OA on the remaining clams (Schwaner et al. 2020). Juvenile clams that had survived the acidified condition for one year appeared to have a broad pH tolerance and performed well under both acidified and control conditions, suggesting they possess a great deal of plasticity. While this could mean there was a selection of animals with this characteristic, it also demonstrates the importance of plasticity under variable pH conditions. The stressful pH conditions were within the range of pH values measured in the environment the parent clams were collected from (Fig. S5). While there were clear phenotypic differences (Schwaner et al. 2020) and a significant regulation of the transcriptome, perhaps a more extreme acidification regime would have induced a higher selective mortality as experimental clams used here appeared to be able to acclimate (i.e., reciprocal transplant) to the lower pH used.

#### Pathways

Overall trends were similar for all life history stages investigated and between RNA and DNA Sequencing. DEGs and SNPs were among genes related to biomineralization, acid–base regulation/ion transport, cytoskeleton, immunity/ stress, and metabolism, highlighting the role these pathways play in resilience to OA.

#### Biomineralization

The impacts on biomineralization are probably the best understood consequences of OA. The upregulation of shell formation genes in this study might represent a mechanistic response to maintain calcification despite OA conditions. Arivalagen et al. (2017) described proteins with domains that are common to the shell matrix of four bivalve species, and they suggested these domains are evolutionary maintained as part of the "basic tool kit" for construction of the CaCO<sub>3</sub> molluscan exoskeleton. These domains included chitin binding-2 and Von Willebrand Factor-A (VWA) (Arivalagan et al. 2017). Chitin binding genes were upregulated under acidification in 2-day, 32 day, and 5 month old clams and a gene with chitin-binding type 2-domain contained a SNP with a higher frequency in acidified clams. Chitin plays a key role in development and functionality in larval bivalve shells (Weiss and Schonitzer 2006). Studies have demonstrated that OA can result in a loss of the quantity of chitin (Mustafa et al. 2015; Zhang et al. 2019), which might explain upregulation of chitin binding genes under OA (Zhao et al. 2020). Another domain of the biomineralization toolkit is VWA, and these domains were found in genes upregulated under OA stress in 2 day, 32 day, and 5 month old clams and a gene with Von Willebrand factor type C domain contained a SNP with higher frequency in acidified clams.

Arivalagan et al. (2017) identified additional proteins (beyond the basic toolkit) that are important for the nucleation and arrangement of  $CaCO_3$  polymorphs, which is particularly interesting because OA causes the dissolution of  $CaCO_3$  polymorphs. These included proteins with epidermal growth factor (EGF), whey acidic protein (WAP), and Fibronectin-3 (FN3) domains (Arivalagan et al. 2017). Genes with EGF domains were upregulated in 2 day, 32 day old, and 5 month old clams under OA stress. SNPs enriched in acidified clams were found in genes containing EGF domains as well. A gene with a WAP domain contained a missense SNP that was enriched in acidified clams and a gene with WAP domain was upregulated in 32 day old postsets. Lastly, c lam larvae upregulated a gene with a FN3 domain.

Aggrecan 1 was upregulated in larval clams raised under OA conditions, and this gene encodes for a calcification related protein similar to perlucin which forms CaCO<sub>3</sub> crystals (Foulon et al. 2019). Receptor-type tyrosine-protein phosphatase had a missense SNP enriched in acidified clams and tyrosine-protein phosphatase is involved in biomineralization (Shi et al. 2012), specifically the formation of the periostracum (Nagai et al. 2007; Zhang et al. 2012) which is a protective layer of shell that is sensitive to OA (Rodolfo-Metalpa et al. 2010; Gazeau et al. 2013; Ramajo et al. 2015). Collagen, type XII, alpha was upregulated in 32 day old postsets from acidified conditions and Collagen alpha-6(VI) chain contained a SNP enriched in acidified clams. Collagens are extracellular matrix proteins that are important for shell matrix formation (Agbaje et al. 2021). Calmodulin has been shown to be important for calcification in the pearl oyster (Yan et al. 2007) and was over expressed under OA in larval sea urchins (Evans et al. 2013) and larval oysters (Dineshram et al. 2015) to sustain calcification under OA. Calmodulin contained a SNP enriched in acidified clams and was upregulated under OA stress at 5 months. Lastly, Vitamin D3 contained a SNP enriched in acidified clams and Vitamin D3 is a stimulator of transepithelial calcium movement during shell mineralization (Lopes-Lima et al. 2012).

#### Acid–Base Regulation/Ion Transport

The supply of calcium and bicarbonate to the site of calcification via transepithelial transport is critical for calcification. Bivalves remove hydrogen ions and transport calcium and bicarbonate ions via membrane bound ion transport proteins (Ramesh et al. 2017; Sillanpaa et al. 2018). Intracellular pH is also regulated through similar mechanisms, making it hard to distinguish between the two processes. The chemistry at the site of calcification dictates the individual response to OA. For example, research has shown that adult bivalves can elevate pH of their calcifying fluids to sustain biomineralization in adverse conditions (Michaelidis et al. 2005). Solute carrier family is a major group of bicarbonate transporters essential to acid-base homeostasis and ion transport in bivalves (Romero et al. 2013). Solute carrier family 26 was identified as an ion transporter involved in providing calcium and bicarbonate for larval calcification (Ramesh et al. 2019). This study showed upregulation of solute carriers in 2 day, 32 day, and 5 month old clams and a solute carrier gene contained a SNP with a higher frequency in clams from low pH conditions. Solute carriers found in this study are part of the major facilitator superfamily (MFS). The MFS has been demonstrated to be involved in acid resistance and intracellular pH homeostasis of Penicillium funiculosum (Xu et al. 2014). Transmembrane transporter activity and transporter activity were enriched GO terms among upregulated DEGs. Upregulating these genes might be a compensatory mechanism to sustain calcification and maintain acid-base regulation despite exposure to low pH external seawater.

#### Cytoskeleton

Similarly, the upregulation of genes related to cytoskeleton could be to maintain cellular integrity and growth in unfavorable conditions. GTP binding protein 6 was upregulated at two days, GTP binding was overexpressed GO term in upregulated genes at 10 months, and GTP binding protein 2 contained a SNP with higher frequency in acidified clams. In a study of larval geoducks, the GO term GTP binding was overexpressed at low pH (Timmins-Schiffman et al. 2019), and c ytoskeletal genes were also overexpressed under OA in the blood clam (Tegillarca granosa; Su et al. 2018). Genes associated with cytoskeleton may also be important for immune function such as cytoskeleton mediated phagocytosis via the actin-myosin contractile system (Allen and Aderem, 1995). In fact, OA was found to decrease the abundance of cytoskeletons in hemocytes from blood clams, resulting in a weakened immune response (Su et al. 2018). Upregulation of cytoskeleton genes could represent a mechanism to increase the ability to produce cytoskeleton components under OA stress.

#### Immunity

Studies have demonstrated that OA can negatively impact bivalve immunity (Bibby et al. 2008; Cao et al. 2018; Su et al. 2018; Schwaner et al. 2020). Baculoviral IAP repeatcontaining protein 2/3, an apoptosis gene, was upregulated under OA stress in 5 month old clams and contained a SNP with a higher frequency in acidified conditions. Ankyrin, which is important to hemocyte function (Wei et al. 2017), contained a SNP enriched in acidified clams. Tripartite motif-containing proteins are involved in pathogen recognition and host defense and are associated with hemocyte function of the scallop *Pecten maximus* (Pauletto et al. 2014). In this study, tripartite motif-containing protein contained a SNP enriched in acidified clams. Similarly, Apolipoprotein B-like contained a SNP with a higher frequency in acidified clams. Apolipoprotein was found in the transcriptome of hemocytes from Manila clam Ruditapes philippinarum and is associated with inflammation (Moreira et al. 2012). Activation of this inflammation pathway is mediated by NFkappa B, and there are several conserved NF-kappa-B sites in the apolipoprotein D promoter. NF- kappa-B actually contained a missense SNP in acidified clams as well. It was downregulated under OA in the blood clam, making them more vulnerable to viral infections as it is important in signaling pathway that induces innate antiviral responses (Liu et al. 2016). Interleukin-1 receptor-associated kinase 4, also associated with hemocytes (Mateo et al. 2012), contained a missense SNP enriched in acidified clams.

In contrast, there were many immune genes downregulated such as peroxidase, a gene with tumor necrosis factor domain, gene with Immunoglobulin domain, and big defensin. Subsets of the same clams used in this study were challenged with pathogenic Vibrio spp. and were found to have a marked reduction in immunity under OA stress (Schwaner et al. 2020). Heat shock protein 70kDA 12A-like was downregulated in 32 day old post set juveniles. While heat shock proteins have been upregulated under OA stress in corals (Moya et al. 2015) and bivalves (Goncalves et al. 2017), studies have also shown downregulation in urchins (Todgham and Hofmann 2009), oysters (Wang et al. 2016; Schwaner et al. 2020), and starfish (Henroth et al. 2011). This might be due to the fact that extended over-expression of heat shock proteins is not sustainable. Reduced expression of heat shock proteins could weaken an organism's ability to mitigate the impacts of OA stress at the cellular level.

#### Metabolism

Other downregulated genes which could demonstrate physiological costs of tolerance to OA were related to metabolism. In 5 and 10 month old juveniles, several metabolic genes displayed decreased expression in acidified conditions. Like with 32 day old clams, 10 month old clams had a general trend of downregulation, which might be due to metabolic depression. Metabolic depression as a mechanism to combat OA has been demonstrated in other marine invertebrate species, including blood clam T. granosa (Zhao et al. 2017), mussels M. chilensis (Navarro et al. 2013) and M. galloprovincialis (Michaelidis et al. 2005). When exposed to acidification, the coral polyp (Acropora millepora; Moya et al. 2012), larval sea urchin (Strongylocentrotus purpuratus; O'Donnell et al. 2010), and larval oyster (C. gigas; Liu et al. 2020) downregulated genes involved in metabolism. Metabolic depression might be an energy saving strategy for surviving under OA conditions. However, this might have associated costs in addition to immune depression. The 10 month old clams downregulated solute carrier and insoluble shell matrix protein, which could come at the cost of shell formation.

#### Conclusions

This study underlines the resilience of *M. mercenaria* to OA and identifies the associated molecular features. As hypothesized, OA appears to significantly influence the transcriptomes of clams and there were DEGs between control and acidified clams at every life stage investigated. While the genetic differentiation between clams surviving 10 months of exposure to OA and control clams was lower than expected, there was convergence of evidence of DEGs and SNPs in similar genes and pathways. Specifically, shell formation, bicarbonate transport, and cytoskeleton were associated with resilience to OA, while reduction in immunity and metabolism might represent costs.

Supplementary Information The online version contains supplementary material available at https://doi.org/10.1007/s10126-022-10183-3.

Acknowledgements The authors would like to thank Drs. Christopher Gobler and Andrew Griffith for their assistance in analyzing DIC samples. We would also like to thank Christopher Paparo at the Southampton Marine Station for assistance with experimental set-up and maintenance.

Author Contributions CS contributed to formal analysis, investigation, methodology, and writing the original draft. SF contributed to formal analysis and manuscript revision. AT and IB assisted with investigation and methodology. MB helped in investigation. EPE helped in methodology, manuscript revision, and funding. BA was integral to funding, investigation, conceptualization, project administration, and reviewing and editing.

Funding This work was supported by funding through the National Oceanic and Atmospheric Administration (NOAA, USA; 67208) Northeast Sea Grant College Consortium in partnership with NOAA Ocean Acidification Program and the New York Sea Grant (NYSG) program to Bassem Allam and Emmanuelle Pales Espinosa (Project R/XG-24). Partial support was also provided by NYSG via Grant # R/XG-32 (East Coast Hard Clam Selective Breeding Collaborative). Additional support was provided by the Mid-Atlantic Sea Grant NOAA Ocean Acidification Program's Ocean Coastal and Estuarine Acidification Graduate Research Fellowship awarded to Caroline Schwaner.

**Data Availability Statement** The original contributions presented in the study are publicly available. Data can be found at: https://www. ncbi.nlm.nih.gov/sra under BioProject PRJNA887696 accession numbers SAMN31181489- SAMN31181584 (RADSeq) and BioProject PRJNA887747 accession numbers SAMN31183563- SAMN31183594 (RNASeq).

#### Declarations

Conflict of Interest The authors declare no conflict of interest.

# References

- Agbaje OBA, Dominguez G, Jabo DE (2021) Organic biopolymers of venus clams: Collagen related matrix in the bivalve shells with crossed lamellar ultrastructure. Biochem Biophys Rep. https:// doi.org/10.1016/j.bbrep.2021.100939
- Allen LH, Aderem A (1995) A role for MARCKS, the alpha isozyme of protein kinase C and myosin I in zymosan phagocytosis by macrophages. J Exp Med 182:182–829
- Arivalagan J, Yarra T, Marie B, Sleight VA, Duvernois-Berthet E, Clark MS, Marie A, Berland S (2017) Insights from the Shell Proteome: Biomineralization to Adaptation. Mol Biol Evol 34:66–77
- Baumann H, Wallace RB, Tagliaferri T, Gobler CJ (2015) Large Natural pH,  $CO_2$ , and  $O_2$  fluctuations in a temperate tidal salt marsh on diel, seasonal, and interannual time scales. Estuaries Coasts 38:220–231
- Bibby R, Widdicombe S, Parry H, Spicer J, Pipe R (2008) Effects of ocean acidification on the immune response of the blue mussel *Mytilus edulis*. Aquat Biol 2:67–74
- Bitter MC, Kapsenber L, Gattuso JP, Pfister CA (2019) Standing genetic variation fuels rapid adaptation to ocean acidification. Nat Commun 10:5821
- Boutet I, Monteiro HJA, Baudry L, Takeuchi T, Bonnivard E, Billoud B, Tanugy A (2022) Chromosomal assembly of the flat oyster (Ostrea edulis L.) genome as a new genetic resource for aquaculture. Evol Appl. https://doi.org/10.1111/eva.13462
- Burge CA, Eakin CM, Friedman CS, Froelich B, Hershverger PK, Hofmann EE, Petes LE, Prager KC, Weil E, Willis BL, Ford SE, Harvell CD (2014) Climate Change influences on Marine Infectious Diseases: Implications for Management and Society. Ann Rev Mar Sci 6:249–277
- Cai WJ, Hu X, Huang WJ, Murrell MC, Lehrter JC, Lohrenz SE, Chou WC, Zhai W, Hollibaugh JT, Wang Y, Zhao P, Guo X, Gunderson K, Dai M, Gong GC (2011) Acidification of subsurface coastal waters enhanced by eutrophication. Nat Geosci 4:766–770
- Cao R, Wang Q, Yang D, Liu Y, Ran W, Qu Y, Zhao J (2018) CO<sub>2</sub>-induced ocean acidification impairs the immune function of the Pacific oyster against *Vibrio splendidus* challenge: An integrated study from a cellular and proteomic perspective. Sci Total Environ 625:1574–1583.
- Cerrato RM, Caron DA, Lonsdale DJ, Rose JM, Schaffner RA (2004) Effect of the northern quahog *Mercenaria mercenaria* on the development of blooms of the brown tide algae *Aureococcus anophagefferens*. MEPS 281:93–108
- Dame RF (2011) Ecology of marine bivalves: an ecosystem approach. CRC Press, Boca Raton
- De Wit P, Durland E, Ventura A, Langdon CJ (2018) Gene expression correlated with delay in shell formation in larval Pacific oysters (Crassostrea gigas) exposed to experimental ocean acidification provides insights into shellformation mechanisms. BMC Genom 19(1):1–15. https://doi.org/10.1186/s12864-018-4519-y
- Dineshram R, Quan Q, Sharma R, Chandramouli K, Yalamanchili HK, Chu I, Thiyagarjan V (2015) Comparative and quantitative proteomics reveal the adaptive strategies of oyster larvae to ocean acidification. Proteomics 15:4120–4134
- Evans TG, Chang F, Menge BA, Hofmann GE (2013) Transcriptomic responses to ocean acidification in larval sea urchins from a naturally variable pH environment. Mol Ecol 22:1609–1625
- Fabry VJ, Seibel BA, Feely RA, Orr JC (2008) Impacts of Ocean Acidification on marine fauna and ecosystem processes. ICES J Mar Sci 65:414–432
- Farhat S, Tanguy A, Pales Espinosa E, Guo X, Boutet I, Smolowitz R, Murphy D, Rivara G, Allam B (2020) Identification of variants associated with hard clam, *Mercenaria mercenaria*, resistance to Quahog Parasite Unknown disease. Genomics 112:4887–4896

- Farhat S, Bonnivard E, Pales Espinosa E, Tanguy A, Boutet I, Guiglielmoni N, Flot JF, Allam B (2022) Comparative analysis of the *Mercenaria mercenaria* genome provides insights into the diversity of transposable elements and immune molecules in bivalve mollusks. BMC Genomics 23:192
- Feely RA, Sabine CL, Lee K, Berelson W, Kleypas J, Fabry VJ, Millero FJ (2004) Impacts of Anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> System in the Oceans. Science 305:362–366
- Feely RA, Sc D, Cooley SR (2009) Ocean acidification: Present conditions and future changes in a high-CO<sub>2</sub> world. Oceanography 22:36-47
- Foulon V, Boudry P, Artiguad S, Guerard F, Hellio C (2019) In Silico Analysis of the Pacific Oyster (*Crassostrea gigas*) Transcriptome over Developmental Stages Reveals Candidate Genes for Larval Settlement. Int J Mol Sci 20:197
- Garcia-Esquivel Z, Bricelj VM, Gonzalez-Gomez MA (2001) Physiological basis for energy demands and early postlarval mortality in the Pacific oyster, *Crassostrea gigas*. J Exp Mar Biol Ecol 263:77–103
- Gattuso J-P, Epitalon J-M, Lavigne H Orr J (2018) seacarb: seawater carbonate chemistry. R package version 3.2.10. http://CRAN.Rproject.org/package=seacarb
- Gazeau FL, Parker LM, Comeau S, Gattuso JP, O'Connor WA, Martin S, Portner HO, Ross PM (2013) Impacts of ocean acidification on marine shelled molluscs. Mar Biol 160:2207–2245
- Goncalves P, Jones DB, Thompson EL, Parker LM, Ross PM, Raftos DA (2017) Transcriptomic profiling of adaptive responses to ocean acidification. Mol Ecol 26:5974–5988
- Gray MW, Langdon CJ, Waldbusser GG, Hales B, Kramer S (2017) Mechanistic understanding of ocean acidification impacts on larval feeding physiology and energy budgets of the mussel *Mytilus californianus*. Mar Ecol Prog Ser. https://doi.org/10.3354/meps11977
- Helm MM, Bourne N, Lovatelli A (2004) Hatchery culture of bivalves: A practical manual. Food and Agriculture Organization of the United Nations, Rome
- Hernroth B, Baden S, Thorndyke M, Dupont S (2011) Immune suppression of the echinoderm *Asterias rubens* (L.) following longterm ocean acidification. Aquat 103:222–224
- Hettinger A, Sanford E, Hill TM, Russell AD, Sato KNS, Hoey J, Forsch M, Page HN, Gaylord B (2012) Persistent carry-over effects of planktonic exposure to ocean acidification in the Olympia oyster. ESA 93:2758–2768
- Lanning G, Eilers S, Pörtner HO, Sokolova IM, Bock C (2010) Impact of Ocean Acidification on Energy Metabolism of Oyster, *Crassostrea gigas*—Changes in Metabolic Pathways and Thermal Response. Mar Drugs 8:2318–2339
- Laurent A, Fennel K, Cai WJ, Huang WJ, Barbero L, Wanninkhof R (2017) Eutrophication-induced acidification of coastal waters in the northern Gulf of Mexico: Insights into origin and processes from a coupled physical-biogeochemical model. Geophys Res Lett 44:946–956
- Li H (2011) A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. 127:2987–03
- Liu S, Shi W, Guo C, Zhao X, Han Y, Peng C, Chai X, Liu G (2016) Ocean acidification weakens the immune response of blood clam through hampering NF-kappa B and toll-like receptor pathways. Fish and Shellfish Immunol 54:322–327
- Liu Z, Zhang Y, Zhou Z, Zong Y, Zheng Y, Liu C, Song L (2020) Metabolomic and transcriptomic profiling reveals the alteration of energy metabolism in oyster larvae during initial shell formation and under experimental ocean acidification. Sci Rep 10:1–11
- Lopes-Lima M, Freitas S, Pereira L, Gouveia E, Hinzmann M, Checa A, Machado J (2012) Ionic regulation and shell mineralization in

the bivalve *Anodonta cygnea* (swan mussel) following heavy-metal exposure. Canadian J Zool 90. https://doi.org/10.1139/z11-129

- López-Landavery EA, Carpizo-Ituarte EJ, Pérez-Carrasco L, Díaz F, Lafarga-De la Cruz F, García-Esquivel Z, Galindo-Sánchez CE (2021) Acidification stress effect on umbonate veliger larval development in *Panopea globosa*. Mar Pollut Bull 163:111945
- Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550
- Mateo DR, Greenwood SJ, Araya MT, Berthe FCJ, Johnson GR, Siah A (2012) Differential gene expression of γ-actin, Tolllike receptor 2 (TLR-2) and interleukin-1 receptor-associated kinase 4 (IRAK-4) in Mya arenaria haemocytes induced by in vivo infections with two Vibrio splendidus strains. Dev Comp Immunol 34:710–714
- Michaelidis B, Ouzounis C, Paleras A, Pörtner HO (2005) Effects of longterm moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. MEPS 293:109–118
- Miller AW, Reynolds AC, Sobrino C, Riedel GF (2009) Shellfish Face Uncertain Future in high CO<sub>2</sub> World: Influence of Acidification on Oyster Larvae Calcification and Growth in Estuaries. PLoS ONE 4(5). https://doi.org/10.1371/journal.pone.0005661
- Millero FJ (2010) Carbonate constants for estuarine waters. Mar Freshw Res 62:139–142
- Moreira R, Balseiro P, Planas JV, Fuste B, Beltran S, Novoa B (2012) Transcriptomics of In Vitro Immune-Stimulated Hemocytes from the Manila Clam *Ruditapes philippinarum* Using High-Throughput Sequencing. PLoS ONE 7(4). https://doi.org/10. 1371/journal.pone.0035009
- Moya A, Huisman L, Ball EE, Hayward DC, Grasso LC, Chua CM, Miller DJ (2012) Whole transcriptome analysis of the coral Acropora millepora reveals complex responses to CO2-driven acidification during the initiation of calcification. Mol Ecol 21:2440–2454
- Moya A, Huisman L, Foret S, Gattuso JP, Hayward DC, Ball EE, Miller DJ (2015) Rapid acclimation of juvenile corals to CO2mediated acidification by upregulation of heat shock protein and Bcl-2 genes. Mol Ecol 24:438–452
- Mustafa S, Kharudin SN, Kian AYS (2015) Effect of Simulated Ocean Acidification on Chitin Content in the Shell of White Shrimp, *Litopenaeus vannamei*. J Fish Sci.com 9(2). https://doi.org/10. 3389/fphys.2021.813582
- Nagai K, Yano M, Morimoto K, Miyamoto H (2007) Tyrosinase localization in mollusc shells. Comp Biochem Physiol B Biochem Mol Biol 146:207–214
- Navarro JM, Torres R, Acuña K, Duarte C, Manriquez PH, Aguilera LM, V, (2013) Impact of medium-term exposure to elevated pCO2 levels on the physiological energetics of the mussel *Mytilus chilensis*. Chemosphere 90:1242–1248
- NMFS National Marine Fisheries Services (2019) Fisheries of the United States 2019. Current Fishery Statistics No. 2019
- O'Donnell MJ, Todgham AE, Swewell MA, Hammond LM, Ruggiero K, Fangue NA, Zippay ML, Hofmann GE (2010) Ocean acidification alters skeletogenesis and gene expression in larval sea urchins. MEPS 398:157–171
- Pauletto M, Milan M, Moreira R, Novoa B, Figueras A, Babbucci M, Patarnello T, Bargelloni. Deep transcriptome sequencing of Pecten maximus hemocytes: A genomic resource for bivalve immunology. Fish and Shellfish Immunol 37:154–165
- Pespeni MH, Sanford E, Gaylord B, Hosfelt HTM, JD, Jaris HK, LaVigne M, Lenz E, Russell AD, Young MK, Palumbi SR, (2013a) Evolutionary change during experimental ocean acidification. PNAS 110:6937–6942
- Pespeni MH, Chang F, MEnge BA, PAlumbi SR, (2013b) Signs of adaptation to local pH conditions across an environmental mosaic in the California current ecosystem. Integr Comp Biol 53:857–870

- Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double Digest RADseq: An inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-Model Species. PLoS ONE 7:e37135
- Ramajo L, Marba N, Prado L, Peron S, Lardies M, Rodriguez-Navarroo AV, Christian A, LAgos NA, Duarte CM, (2015) Biomineralization changes with food supply confer juvenile scallops (*Agropecten purpuratus*) resistance to ocean acidification. Glob Change Biol 22:2025–2037
- Ramesh K, Hu MY, Thomsen J, Bleich M, Melzner F (2017) Mussel larvae modify calcifying fluid carbonate chemistry to promote calcification. Nat Commun 8:1709
- Ramesh K, Yarra T, Clark MS, John U, Melzner F (2019) Expression of calcification-related ion transporters during blue mussel larval development. Ecol Evol 9:77157–77172
- Rochette NC, Rivera-Colón AG, Catchen JM (2019) Stacks 2: Analytical methods for paired-end sequencing improve RADseq-based population genomics. Mol Ecol 28:4737–4754
- Rodolfo-Metalpa R, Lombardi C, Cocito S, Hall-Spencer JM, Gambi MC (2010) Effects of ocean acidification and high temperatures on the bryozoan *Myriapora truncata* at natural CO<sub>2</sub> vents. Mar Ecol 31:447–456
- Romero MF, Chen AP, Parker MD, Boron WF (2013) The SLC4 family of bicarbonate (HCO<sub>3</sub><sup>-</sup>) transporters. Mol Aspects Med 34:159–182
- Ropp AJ (2020) Population Structure of the Hard Clam, *Mercenaria mercenaria*, along the EAst Coast of North America. Dissertations, Theses, and Masters Projects. Paper 1616444311. https:// doi.org/10.25773/v5-wa9q-8c19
- Rice MA (1992) The Northern Quahog: The biology of *Mercenaria mercenaria*. Rhode Island Sea Grant ISBN:0–938412–33–7
- Sabine CL, Feely RA, Gruber N, Key RM, Lee K, Bullister JL, WAnninkhof R, Wong CS, Wallace DWR, Tilbrook B, Millero FJ, Peng TH, Kozyr A, Ono T, Rios AF, (2004) The Ocean Sink for Anthropogenic CO<sub>2</sub>. Science 305:367–371
- Schwaner C, Barbosa M, Connors P, Park TJ, de Silva D, Griffith A, Gobler CJ, Espinosa EP, Allam B (2020) Experimental acidification increases susceptibility of *Mercenaria mercenaria* to infection by *Vibrio* species. Mar Environ Res 154:104872
- Shi Y, Yu C, Gu Z, Zhan X, Wang Y, Wang A (2012) Characterization of the Pearl Oyster (*Pinctada martensii*) Mantle Transcriptome Unravels Biomineralization Genes. Mar Biotechnol 15:175–187
- Sillanpaa KJ, Sundh H, Sundell KS (2018) Calcium transfer across the outer mantle epithelium in the Pacific oyster. Crassostrea Gigas Proc Royal Soc b 285:20181676
- Stony Brook University (2016) Understanding SeaWulf. https://it. stonybrook.edu/help/kb/understanding-seawulf
- Su W, Rong J, Zha S, Yan M, Fang J, Liu G (2018) Ocean acidification affects the cytoskeleton, lysozymes, and nitric oxide of hemocytes: a possible explanation for the hampered phagocytosis in blood clams. Tegillarca Granosa Front Physiol 9:619
- Sun T, Tang X, Jiang Y, Wang Y (2017) Seawater acidification induced immune function changes of haemocytes in *Mytilus edulis*: a comparative study of CO<sub>2</sub> and HCl enrichment. Sci Rep 7:41488
- Talmage SC, Gobler CJ (2009) The effects of elevated carbon dioxide concentrations on the metamorphosis, size, and survival of larval hard clams (*Mercenaria mercenaria*), bay scallops (*Argopecten irradians*), and Eastern oysters (*Crassostrea virginica*). Limnol Oceangr 54:2072–2080
- Talmage SC, Gobler CJ (2010) Effects of past, present, and future ocean carbon dioxide concentrations on the growth and survival or larval shellfish. PNAS 107:17246–17251
- Talmage SC, Gobler CJ (2011) Effects of elevated temperature and carbon dioxide on the Growth and Survival of Larvae and Juveniles of Three Species of Northwest Atlantic Bivalves. PLOS ONE 6(10). https://doi.org/10.1371/journal.pone.0026941

- Thomsen J, Melzner F (2010) Moderate seawater acidification does not elicit long-term metabolic depression in the blue mussel *Mytilus edulis*. Mar Biol 157:2667–2676
- Thomsen J, Stapp LS, Haynert K, Schade H, Danelli M, Lanning G, Wegner KM, Melzner F (2017) Naturally acidified habitat selects for ocean acidification-tolerant mussels. Sci Adv. https://doi.org/ 10.1126/sciadv.1602411
- Timmins-Schiffman, O'Donnell MJ, Friedman CS, Roberts SB (2012) Elevated pCO<sub>2</sub> causes developmental delay in early larval Pacific oysters, *Crassostrea gigas*. Mar Bio 160:1973–1982
- Timmins-Schiffman E, Coffey WD, Hua W, Nunn BL, Dickinson GH, Roberts SB (2014) Shotgun proteomics reveals physiological response to ocean acidification in *Crassotrea gigas*. BMC Genomics 15:951
- Timmins-Schiffman E, Guzman JM, Elliott R, Vadopalas B, Roberts SB (2019) Dynamic response in the larval geoduck clam proteome to elevated *p*CO<sub>2</sub>. BioRxiv. https://doi.org/10.1101/613018
- Todgham AE, Hofmann GE (2009) Transcriptomic response of sea urchin larvae *Strongylocentrotus purpuratus* to CO2-driven seawater acidification. J Exp Biol 212:2579–2594
- Uthicke S, Deshpande NP, Liddy M, Patel F, Lamare M, Wilkins MR (2019) Little evidence of adaptation potential to ocean acidification in sea urchins living in "Future Ocean" conditions at a CO<sub>2</sub> vent. Ecol Evol 9:10004–10016
- Videla JA, Chaparro OR, Thompson RJ, Concha II (1998) Role of biochemical energy reserves in the metamorphosis and early juvenile development of the oyster Ostrea chilensis. Mar Biol 132:635–640
- Waldbusser GG, Brunner EL, Haley BA, Hales B, Langdon CJ, Prahl FG (2013) A developmental and energetic basis linking larval oyster shell formation to acidification sensitivity. Geophy Res Lett 40:2171–2176
- Waldbusser GG, Hales B, Langdon CJ, Haley BA, Schrader P, Brunner EL, Hutchinson G (2015) Ocean acidification has multiple modes of action on bivalve larvae. PLoS ONE 10:e0128376
- Wall CC, Peterson BJ, Gobler CJ (2008) Facilitation of seagrass Zostera marina productivity by suspension-feeding bivalves. MEPS 357:165–174
- Wallace RK, Waters P, Rikard RS (2008) Oyster hatchery techniques. Southern Regional Aquaculture Center
- Wallace RB, Baumann H, Grear JS, Aller RC, Gobler CJ (2014) Coastal Ocean Acidification: The other eutrophication problem. Estuar Coast Shelf Sci 148:1–13
- Wang Q, Cao R, Ning X, You L, Mu C, Wang C, Wei L, Cong M, Wu H, Zhao J (2016) Effects of ocean acidification on immune response of the Pacific oyster *Crassostrea gigas*. Fish Shellfish Immunol 49:24–33
- Wei J, Liu B, Fan S, Li H, Chen M, Zhang B, Yu D (2017) Differentially expressed immune-related genes in hemocytes of the pearl oyster *Pinctada fucata* against allograft identified by transcriptome analysis. Fish & Shellfish Immunol 62:247–256

- Weiss IM, Schonitzer V (2006) The distribution of chitin in larval shells of the bivalve mollusk *Mytilus galloprovincialis*. J Struct Biol 153:264–277
- Wessel N, Martin S, Badou A, Dubois P, Huchette S, Julia V, Nunes F, Harney E, Paillard C, Auzoux-Bordenave S (2018) Effect of CO2induced ocean acidification on the early development and shell mineralization of the European abalone (*Haliotis tuberculata*). J Exp Mar Biol Ecol 508:52–63
- White MM, McCorkle DC, Mullineaux LS, Cohen AL (2013) Early Exposure of Bay Scallops (*Argopectin irradians*) to High CO2 Causes a Decrease in Larval Shell Growth. PLos ONE 8(4). https://doi.org/10. 1371/journal.pone.0061065
- Xu X, Chen J, Xu H, Li D (2014) Role of a major facilitator superfamily transporter in adaptation capacity of *Penicillium funiculosum* under extreme acidic stress. Fungal Genet Biol 69:75–83
- Xu X, Yang F, Zhao L, Yan X (2016) Seawater acidification affects the physiological energetics and spawning capacity of the Manila clam *Ruditapes philippinarum* during gonadal maturation. Comp. Biochem Physiol Part A Mol Integr Physiol 196:20–29
- Yan Z, Fang Z, Ma Z, Deng J, Li S, Xie L, Zhang R (2007) Biomineralization: functions of calmodulin-like protein in the shell formation of pearl oyster. Biochim Biophys Acta 1170:1338–1344
- Zarnoch CB, Schreibman MP (2008) Influence of temperature and food availability on the biochemical composition and mortality of juvenile *Mercenaria mercenaria* (L.) during the over-winter period. Aquacul 274:281–291
- Zhang G, Fang X, Guo X, Li L, Luo R, Xu F, Yang P, Zhang L, Wang J (2012) The oyster genome reveals stress adaptation and complexity of shell formation. Nature 490:49–54
- Zhang Y, Liu Z, Song X, Huang S, Wang L, Song L (2019) The Inhibition of Ocean Acidification on the Formation of Oyster Calcified Shell by Regulating the Expression of Cg chs1 and Cg chit4. Front Physiol 10:1034
- Zhao X, Shi W, Han Y, Liu S, Guo C, Fu W, Liu G (2017) Ocean acidification adversely influences metabolism, extracellular pH and calcification of an economically important marine bivalve, *Tegillarca granosa*. Mar Environ Res 125:82–89
- Zhao X, Han Y, Chen B, Xia B, Qu K, Liu G (2020) CO<sub>2</sub>-driven ocean acidification weakens mussel shell defense capacity and induces global molecular compensatory responses. Chemosphere 243:125415

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.